

## General Information

Amino Acid Chelated **INTRACEPT™** is manufactured and designed as a foliar nutrient. **INTRACEPT™** is designed for use on large variety of crops, including vegetables, tree fruits, small fruits, citrus trees and ornamentals. **INTRACEPT™** should be applied at planting or at the beginning of stressful growth stages such as first true leaf, flowering, and fruit set. **INTRACEPT™** is readily available and rapidly absorbed by the plant because of the natural complexing and chelation properties. It is designed for foliar, row, or broadcast application, and it is water soluble and nontoxic to plants when applied as directed.

**INTRACEPT™** is a proprietary formulation and blend of plant extracts, natural polysaccharides, and Amino Acid Chelated micronutrients. This product has been designed to promote nutrient penetration, uptake, and utilization. **INTRACEPT™** has also been specially reacted with emulsion polymerization that provides a wide range of mixing ease with other fertilizers or chemicals.

## Guaranteed Analysis

Sulfur (S) .....	1.0%
Copper (Cu) .....	0.1%
Iron (Fe) .....	0.1%
Manganese (Mn) .....	0.33%
Zinc (Zn) .....	0.4%

*(Derived from copper sulfate, ferrous sulfate, manganese sulfate, zinc sulfate, chelated with Amino Acids.)*

### ALSO CONTAINS NONPLANT FOOD INGREDIENTS

Polysaccharides .....	0.82%
Organic Acids .....	1.0%

*(Derived from plant extracts and fermentation, amino acids derived from kelp)*

### Product Specifications

Amino Acid chelated **INTRACEPT™** is a yield enhancing formulation that contains biostimulants and Amino Acid chelated trace minerals. **INTRACEPT™** enhances plant growth and development by stimulating nutrient uptake, nutrient penetration, and nutrient utilization.

Manufactured by  
**Northwest Agricultural Products, Inc.**  
 821 S. Chestnut • Pasco WA 99301  
**509 / 547-8234**

© 2003 NAP 09/2007 QA# 0442 m/s

# INTRACEPT™

## A Chelated Micronutrient & Plant Growth Enhancer



Distributed by Northwest Agricultural Products  
 P.O. Box 3453 - Pasco, WA 99302  
**509 / 547-8234**

Net Contents: 5 gallons   
 2.5 gallons

Weight per Gallon: 9.75 lbs.



Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.htm>

**SHAKE WELL BEFORE USING, DO NOT FREEZE**  
**CAUTION: KEEP OUT OF THE REACH OF CHILDREN**  
**HARMFUL IF SWALLOWED**

## Directions for Use

### FOLIAR APPLICATION

Three weeks after emergence (or as otherwise recommended by a certified crop advisor) apply recommended rates of **INTRACEPT™**. This product may be mixed with pesticides and fertilizers. Three or four weeks after the first application (or as recommended by a certified crop advisor), apply the recommended rate of **INTRACEPT™**. Seven to ten days after the second application (or as otherwise recommended by a certified crop advisor) apply the recommended rate of **INTRACEPT™**.

Dilute one part **INTRACEPT™** with 20 or more parts water. Apply by a spray method and in an adequate amount of water that will provide complete coverage of the plants. Use of a non-ionic wetting agent may improve spray coverage of certain hard to wet plants. Do Not Apply Undiluted. **INTRACEPT™** can be included in a regular spray program on crops. Consult with a Northwest Ag Products representative on compatibility with other spray materials. The rate of application will be depend on the crop, stage of growth, and severity of deficiency. The maximum recommended rates are for mature, full sized plants. Reduce the rates proportionately when spraying smaller plants.

### Recommendations

**INTRACEPT™** may be used on legumes, grain, crops, root crops, cucurbits, cole crops, leafy vegetables, woody and herbaceous ornamentals, deciduous fruits, vine crops, tropical and subtropical fruits, and many other crops.

**Alfalfa:** Apply 2 to 32 fl. oz. per acre during periods of first growth and after first cutting at 4th to 6th leaf stage. The applications may be repeated 2 or more times through the growing season.

**Wheat:** Apply 2 to 32 fl. oz. per acre when plants are 6" high and 12 to 14 days before booting.

**Beans:** Apply 2 to 32 fl. oz. per acre at first cultivation and 12 to 14 days before blossoming.

**Potatoes:** Apply 16 to 32 fl. oz. per acre at row closure. Application should be applied during periods of rapid growth or nutritional stress. Applications may be repeated 2 or more times through growing season.

**Rice:** Apply 2 to 32 fl. oz. per acre just before booting.

**Fruit Trees:** Apply 2 to 32 fl. oz. per acre before blossoming and shortly after petal fall.

**Melons and Vine Crops:** Apply 2 to 32 fl. oz. per acre at first female flower and again at softball size fruit.

**Onions:** Apply 2 to 32 fl. oz. per acre at 4 to 6 leaf stage.

### PRECAUTIONARY STATEMENTS

In all cases, it is recommended that a small scale test for compatibility to be conducted before mixing a full batch.

### CONDITION OF SALE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. Since weather, crop, soil, and other conditions may vary, NAP and/or seller make no warranty of any kind, expressed or implied, concerning the use of this product. The user assumes all risks of use of handling, whether or not in accordance with directions or suggestions.